

RESEARCH

Open Access



Approximation of a kind of new type Bézier operators

Mei-Ying Ren^{1*} and Xiao-Ming Zeng²

*Correspondence:

npmeiyingr@163.com

¹School of Mathematics and
Computer Science, Wuyi University,
Wuyishan, 354300, China
Full list of author information is
available at the end of the article**Abstract**

In this paper, a kind of new type Bézier operators is introduced. The Korovkin type approximation theorem of these operators is investigated. The rates of convergence of these operators are studied by means of modulus of continuity. Then, by using the Ditzian-Totik modulus of smoothness, a direct theorem concerned with an approximation for these operators is also obtained.

MSC: 41A10; 41A25; 41A36**Keywords:** Bézier type operators; Korovich type approximation theorem; rate of convergence; direct theorem; modulus of smoothness

1 Introduction

In view of the Bézier basis function, which was introduced by Bézier [1], in 1983, Chang [2] defined the generalized Bernstein-Bézier polynomials for any $\alpha > 0$, and a function f defined on $[0, 1]$ as follows:

$$B_{n,\alpha}(f; x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) [J_{n,k}^\alpha(x) - J_{n,k+1}^\alpha(x)], \quad (1)$$

where $J_{n,n+1}(x) = 0$, and $J_{n,k}(x) = \sum_{i=k}^n P_{n,i}(x)$, $k = 0, 1, \dots, n$, $P_{n,i}(x) = \binom{n}{i} x^i (1-x)^{n-i}$. $J_{n,k}(x)$ is the Bézier basis function of degree n .

Obviously, when $\alpha = 1$, $B_{n,\alpha}(f; x)$ become the well-known Bernstein polynomials $B_n(f; x)$, and for any $x \in [0, 1]$, we have $1 = J_{n,0}(x) > J_{n,1}(x) > \dots > J_{n,n}(x) = x^n$, $J_{n,k}(x) - J_{n,k+1}(x) = P_{n,k}(x)$.

During the last ten years, the Bézier basis function was extensively used for constructing various generalizations of many classical approximation processes. Some Bézier type operators, which are based on the Bézier basis function, have been introduced and studied (e.g., see [3–9]).

In 2012, Ren [10] introduced Bernstein type operators as follows:

$$L_n(f; x) = f(0)P_{n,0}(x) + \sum_{k=1}^{n-1} P_{n,k}(x)B_{n,k}(f) + f(1)P_{n,n}(x), \quad (2)$$

where $f \in C[0, 1]$, $x \in [0, 1]$, $P_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$, $k = 0, 1, \dots, n$, and $B_{n,k}(f) = \frac{1}{B(nk, n(n-k))} \int_0^1 t^{nk-1} (1-t)^{n(n-k)-1} f(t) dt$, $k = 1, \dots, n-1$, $B(\cdot, \cdot)$ is the beta function.

The moments of the operators $L_n(f; x)$ were obtained as follows (see [10]).

Remark 1 For $L_n(t^j; x)$, $j = 0, 1, 2$, we have

- (i) $L_n(1; x) = 1$;
- (ii) $L_n(t; x) = x$;
- (iii) $L_n(t^2; x) = \frac{n(n-1)}{n^2+1}x^2 + \frac{n+1}{n^2+1}x$.

In the present paper, we will study the Bézier variant of the Bernstein type operators $L_n(f; x)$, which have been given by (2). We introduce a new type of Bézier operators as follows:

$$L_{n,\alpha}(f; x) = f(0)Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x)B_{n,k}(f) + f(1)Q_{n,n}^{(\alpha)}(x), \quad (3)$$

where $f \in C[0, 1]$, $x \in [0, 1]$, $\alpha > 0$, $Q_{n,k}^{(\alpha)}(x) = J_{n,k}^\alpha(x) - J_{n,k+1}^\alpha(x)$, $J_{n,n+1}(x) = 0$, $J_{n,k}(x) = \sum_{i=k}^n P_{n,i}(x)$, $k = 0, 1, \dots, n$, $P_{n,i}(x) = \binom{n}{i} x^i (1-x)^{n-i}$, and $B_{n,k}(f) = \frac{1}{B(nk, n(n-k))} \int_0^1 t^{nk-1} (1-t)^{n(n-k)-1} f(t) dt$, $k = 1, \dots, n-1$, $B(\cdot, \cdot)$ is the beta function.

It is clear that $L_{n,\alpha}(f; x)$ are linear and positive on $C[0, 1]$. When $\alpha = 1$, $L_{n,\alpha}(f; x)$ become the operators $L_n(f; x)$.

The goal of this paper is to study the approximation properties of these operators with the help of the Korovkin type approximation theorem. We also estimate the rates of convergence of these operators by using a modulus of continuity. Then we obtain the direct theorem concerned with an approximation for these operators by means of the Ditzian-Totik modulus of smoothness.

In the paper, for $f \in C[0, 1]$, we denote $\|f\| = \max\{|f(x)| : x \in [0, 1]\}$. $\omega(f, \delta)$ ($\delta > 0$) denotes the usual modulus of continuity of $f \in C[0, 1]$.

2 Some lemmas

Now, we give some lemmas, which are necessary to prove our results.

Lemma 1 (see [2]) *Let $\alpha > 0$. We have*

- (i) $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n J_{n,k}^\alpha(x) = x$ uniformly on $[0, 1]$;
- (ii) $\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{k=1}^n k J_{n,k}^\alpha(x) = \frac{x^2}{2}$ uniformly on $[0, 1]$.

Lemma 2 *Let $\alpha > 0$. We have*

- (i) $L_{n,\alpha}(1; x) = 1$;
- (ii) $\lim_{n \rightarrow \infty} L_{n,\alpha}(t; x) = x$ uniformly on $[0, 1]$;
- (iii) $\lim_{n \rightarrow \infty} L_{n,\alpha}(t^2; x) = x^2$ uniformly on $[0, 1]$.

Proof By simple calculation, we obtain $B_{n,k}(1) = 1$, $B_{n,k}(t) = \frac{k}{n}$, $B_{n,k}(t^2) = \frac{1}{n^2+1}(k^2 + \frac{k}{n})$.

(i) Since $\sum_{k=0}^n Q_{n,k}^{(\alpha)}(x) = 1$, by (3) we can get $L_{n,\alpha}(1; x) = 1$.

(ii) By (3), we have

$$\begin{aligned} L_{n,\alpha}(t; x) &= \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) \frac{k}{n} + Q_{n,n}^{(\alpha)}(x) \\ &= [J_{n,1}^{\alpha}(x) - J_{n,2}^{\alpha}(x)] \frac{1}{n} + \cdots + [J_{n,n-1}^{\alpha}(x) - J_{n,n}^{\alpha}(x)] \frac{n-1}{n} + J_{n,n}^{\alpha}(x) \frac{n}{n} \\ &= \frac{1}{n} \sum_{k=1}^n J_{n,k}^{\alpha}(x), \end{aligned}$$

thus, by Lemma 1(i), we have $\lim_{n \rightarrow \infty} L_{n,\alpha}(t; x) = x$ uniformly on $[0, 1]$.

(iii) By (3), we have

$$\begin{aligned} L_{n,\alpha}(t^2; x) &= \frac{1}{n^2+1} \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) \left(k^2 + \frac{k}{n}\right) + Q_{n,n}^{(\alpha)}(x) \\ &= \frac{1}{n^2+1} \sum_{k=1}^n \left(2k - 1 + \frac{1}{n}\right) J_{n,k}^{\alpha}(x) \\ &= \frac{1}{n^2+1} \left[2n^2 \cdot \frac{1}{n^2} \sum_{k=1}^n k J_{n,k}^{\alpha}(x) - n \cdot \frac{1}{n} \sum_{k=1}^n J_{n,k}^{\alpha}(x) + \frac{1}{n} \sum_{k=1}^n J_{n,k}^{\alpha}(x) \right], \end{aligned}$$

thus, by Lemma 1, we have $\lim_{n \rightarrow \infty} L_{n,\alpha}(t^2; x) = x^2$ uniformly on $[0, 1]$. \square

Lemma 3 (see [11]) For $x \in [0, 1]$, $k = 0, 1, \dots, n$, we have

$$0 \leq Q_{n,k}^{(\alpha)}(x) \leq \begin{cases} \alpha P_{n,k}(x), & \alpha \geq 1; \\ P_{n,k}^{\alpha}(x), & 0 < \alpha < 1. \end{cases}$$

Lemma 4 (see [12]) For $0 < \alpha < 1$, $\beta > 0$, we have

$$\sum_{k=0}^n |k - nx|^{\beta} P_{n,k}^{\alpha}(x) \leq (n+1)^{1-\alpha} (A_{\frac{\beta}{\alpha}})^{\alpha} n^{\frac{\beta}{2}},$$

where the constant A_s only depends on s .

Lemma 5 For $\alpha \geq 1$, we have

$$\begin{aligned} \text{(i)} \quad L_{n,\alpha}((t-x)^2; x) &\leq \frac{\alpha}{2} \cdot \frac{1}{n}; \\ \text{(ii)} \quad L_{n,\alpha}(|t-x|; x) &\leq \sqrt{\frac{\alpha}{2}} \cdot \sqrt{\frac{1}{n}}. \end{aligned}$$

Proof (i) By (3), Lemma 3 and Remark 1, we obtain

$$\begin{aligned} L_{n,\alpha}((t-x)^2; x) &= x^2 Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}((t-x)^2) + (1-x)^2 Q_{n,n}^{(\alpha)}(x) \end{aligned}$$

$$\begin{aligned}
&\leq \alpha \left[x^2 P_{n,0}(x) + \sum_{k=1}^{n-1} P_{n,k}(x) B_{n,k}((t-x)^2) + (1-x)^2 P_{n,n}(x) \right] \\
&= \alpha L_n((t-x)^2; x) \\
&= \frac{(n+1)\alpha}{n^2+1} x(1-x),
\end{aligned} \tag{4}$$

since $\max_{0 \leq x \leq 1} x(1-x) = \frac{1}{4}$, and for any $n \in \mathbb{N}$, one can get $\frac{n(n+1)}{n^2+1} \leq 2$, so we have

$$L_{n,\alpha}((t-x)^2; x) \leq \frac{\alpha}{2} \cdot \frac{1}{n}.$$

(ii) In view of $L_{n,\alpha}(1; x) = 1$, by the Cauchy-Schwarz inequality, we have

$$L_{n,\alpha}(|t-x|; x) \leq \sqrt{L_{n,\alpha}(1; x)} \sqrt{L_{n,\alpha}((t-x)^2; x)},$$

thus, we get

$$L_{n,\alpha}(|t-x|; x) \leq \sqrt{\frac{\alpha}{2}} \cdot \sqrt{\frac{1}{n}}.$$

□

Lemma 6 For $0 < \alpha < 1$, we have

- (i) $L_{n,\alpha}((t-x)^2; x) \leq M_\alpha n^{-\alpha};$
- (ii) $L_{n,\alpha}(|t-x|; x) \leq \sqrt{M_\alpha} \cdot n^{-\frac{\alpha}{2}}.$

Here the constant M_α only depends on α .

Proof (i) By (3) and Lemma 3, we obtain

$$\begin{aligned}
&L_{n,\alpha}((t-x)^2; x) \\
&= x^2 Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}((t-x)^2) + (1-x)^2 Q_{n,n}^{(\alpha)}(x) \\
&\leq x^2 P_{n,0}^\alpha(x) + \sum_{k=1}^{n-1} P_{n,k}^\alpha(x) B_{n,k}((t-x)^2) + (1-x)^2 P_{n,n}^\alpha(x) \\
&= \sum_{k=0}^n P_{n,k}^\alpha(x) \left[\frac{1}{n^2+1} \left(k^2 + \frac{k}{n} \right) - 2x \frac{k}{n} + x^2 \right] \\
&= \frac{1}{n^2+1} \sum_{k=0}^n (k-nx)^2 P_{n,k}^\alpha(x) + \frac{1}{n^2+1} \sum_{k=0}^n P_{n,k}^\alpha(x) \left(\frac{k}{n} - 2x \frac{k}{n} + x^2 \right) \\
&:= I_1 + I_2.
\end{aligned}$$

By Lemma 4, we have $I_1 \leq \frac{n(n+1)}{n^2+1} (n+1)^{-\alpha} (A_{\frac{2}{\alpha}})^\alpha \leq 2(A_{\frac{2}{\alpha}})^\alpha n^{-\alpha}$, where the constant $A_{\frac{2}{\alpha}}$ only depends on α .

Using the Hölder inequality, we have $\sum_{k=0}^n P_{n,k}^\alpha(x) \leq (n+1)^{1-\alpha} [\sum_{k=0}^n P_{n,k}(x)]^\alpha$, and $|\frac{k}{n} - 2x\frac{k}{n} + x^2| \leq 4$, so we have

$$\begin{aligned} I_2 &\leq \frac{4}{n^2+1} (n+1)^{1-\alpha} \left[\sum_{k=0}^n P_{n,k}(x) \right]^\alpha \\ &= \frac{4}{n^2+1} (n+1)^{1-\alpha} \leq 4n^{-\alpha}. \end{aligned}$$

Denote $M_\alpha = 2(A_{\frac{2}{\alpha}})^\alpha + 4$, then we can get

$$L_{n,\alpha}((t-x)^2; x) \leq M_\alpha n^{-\alpha}.$$

(ii) Since

$$L_{n,\alpha}(|t-x|; x) \leq \sqrt{L_{n,\alpha}(1; x)} \sqrt{L_{n,\alpha}((t-x)^2; x)},$$

thus, we get

$$L_{n,\alpha}(|t-x|; x) \leq \sqrt{M_\alpha} \cdot n^{-\frac{\alpha}{2}}.$$

□

Lemma 7 For $f \in C[0,1]$, $x \in [0,1]$ and $\alpha > 0$, we have

$$|L_{n,\alpha}(f; x)| \leq \|f\|.$$

Proof By (3) and Lemma 2(i), we have

$$|L_{n,\alpha}(f; x)| \leq \|f\| L_{n,\alpha}(1; x) = \|f\|.$$

□

3 Main results

First of all we give the following convergence theorem for the sequence $\{L_{n,\alpha}(f; x)\}$.

Theorem 1 Let $\alpha > 0$. Then the sequence $\{L_{n,\alpha}(f; x)\}$ converges to f uniformly on $[0,1]$ for any $f \in C[0,1]$.

Proof Since $L_{n,\alpha}(f; x)$ is bounded and positive on $C[0,1]$, and by Lemma 2, we have $\lim_{n \rightarrow \infty} \|L_{n,\alpha}(e_j; \cdot) - e_j\| = 0$ for $e_j(t) = t^j$, $j = 0, 1, 2$. So, according to the well-known Bohman-Korovkin theorem ([13], p.40, Theorem 1.9), we see that the sequence $\{L_{n,\alpha}(f; x)\}$ converges to f uniformly on $[0,1]$ for any $f \in C[0,1]$. □

Next we estimate the rates of convergence of the sequence $\{L_{n,\alpha}\}$ by means of the modulus of continuity.

Theorem 2 Let $f \in C[0,1]$, $x \in [0,1]$. Then

- (i) when $\alpha \geq 1$, we have $\|L_{n,\alpha}(f; \cdot) - f\| \leq (1 + \sqrt{\frac{\alpha}{2}}) \omega(f, \frac{1}{\sqrt{n}})$;
- (ii) when $0 < \alpha < 1$, we have $\|L_{n,\alpha}(f; \cdot) - f\| \leq (1 + \sqrt{M_\alpha}) \omega(f, n^{-\frac{\alpha}{2}})$.

Here the constant M_α only depends on α .

Proof (i) When $\alpha \geq 1$, by Lemma 2(i), we have

$$\begin{aligned}
 & |L_{n,\alpha}(f; x) - f(x)| \\
 & \leq |f(0) - f(x)| Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}(|f(t) - f(x)|) + |f(1) - f(x)| Q_{n,n}^{(\alpha)}(x) \\
 & \leq \omega(f, |0 - x|) Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}(\omega(f, |t - x|)) + \omega(f, |1 - x|) Q_{n,n}^{(\alpha)}(x) \\
 & \leq (1 + \sqrt{n}|0 - x|) \omega\left(f, \frac{1}{\sqrt{n}}\right) Q_{n,0}^{(\alpha)}(x) \\
 & \quad + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}\left((1 + \sqrt{n}|t - x|) \omega\left(f, \frac{1}{\sqrt{n}}\right)\right) \\
 & \quad + (1 + \sqrt{n}|1 - x|) \omega\left(f, \frac{1}{\sqrt{n}}\right) Q_{n,n}^{(\alpha)}(x) \\
 & \leq \omega\left(f, \frac{1}{\sqrt{n}}\right) + \sqrt{n} \omega\left(f, \frac{1}{\sqrt{n}}\right) L_{n,\alpha}(|t - x|; x),
 \end{aligned}$$

so, by Lemma 5(ii), we obtain $|L_{n,\alpha}(f; x) - f(x)| \leq (1 + \sqrt{\frac{\alpha}{2}}) \omega(f, \frac{1}{\sqrt{n}})$. The desired result follows immediately.

(ii) When $0 < \alpha < 1$, by Lemma 2(i), we have

$$\begin{aligned}
 & |L_{n,\alpha}(f; x) - f(x)| \\
 & \leq \omega(f, |0 - x|) Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}(\omega(f, |t - x|)) + \omega(f, |1 - x|) Q_{n,n}^{(\alpha)}(x) \\
 & \leq (1 + n^{\frac{\alpha}{2}} |0 - x|) \omega(f, n^{-\frac{\alpha}{2}}) Q_{n,0}^{(\alpha)}(x) + \sum_{k=1}^{n-1} Q_{n,k}^{(\alpha)}(x) B_{n,k}(1 + n^{\frac{\alpha}{2}} |t - x|) \omega(f, n^{-\frac{\alpha}{2}}) \\
 & \quad + (1 + n^{\frac{\alpha}{2}} |1 - x|) \omega(f, n^{-\frac{\alpha}{2}}) Q_{n,n}^{(\alpha)}(x) \\
 & = \omega(f, n^{-\frac{\alpha}{2}}) + n^{\frac{\alpha}{2}} \omega(f, n^{-\frac{\alpha}{2}}) L_{n,\alpha}(|t - x|; x),
 \end{aligned}$$

so, by Lemma 6(ii), we obtain $|L_{n,\alpha}(f; x) - f(x)| \leq (1 + \sqrt{M_\alpha}) \omega(f, n^{-\frac{\alpha}{2}})$. The desired result follows immediately. \square

Theorem 3 Let $f \in C^1[0, 1]$, $x \in [0, 1]$. Then

(i) when $\alpha \geq 1$, we have

$$|L_{n,\alpha}(f; x) - f(x)| \leq \|f'\| \sqrt{\frac{\alpha}{2n}} + \omega\left(f', \frac{1}{\sqrt{n}}\right) \left(1 + \sqrt{\frac{\alpha}{2}}\right) \sqrt{\frac{\alpha}{2n}};$$

(ii) when $0 < \alpha < 1$, we have

$$|L_{n,\alpha}(f; x) - f(x)| \leq \|f'\| \sqrt{M_\alpha n^{-\alpha}} + \omega\left(f', n^{-\frac{\alpha}{2}}\right) (1 + \sqrt{M_\alpha}) \sqrt{M_\alpha n^{-\alpha}},$$

where the constant M_α only depends on α .

Proof Let $f \in C^1[0, 1]$. For any $t, x \in [0, 1]$, $\delta > 0$, we have

$$\begin{aligned} |f(t) - f(x) - f'(x)(t - x)| &\leq \left| \int_x^t |f'(u) - f'(x)| du \right| \\ &\leq \omega(f', |t - x|)|t - x| \\ &\leq \omega(f', \delta)(|t - x| + \delta^{-1}(t - x)^2), \end{aligned}$$

hence, by the Cauchy-Schwarz inequality, we have

$$\begin{aligned} |L_{n,\alpha}(f(t) - f(x) - f'(x)(t - x); x)| \\ \leq \omega(f', \delta)(L_{n,\alpha}(|t - x|; x) + \delta^{-1}L_{n,\alpha}((t - x)^2; x)) \\ \leq \omega(f', \delta)(\sqrt{L_{n,\alpha}(1; x)} \\ + \delta^{-1}\sqrt{L_{n,\alpha}((t - x)^2; x)})\sqrt{L_{n,\alpha}((t - x)^2; x)}. \end{aligned}$$

So we get

$$\begin{aligned} |L_{n,\alpha}(f; x) - f(x)| \\ \leq \|f'\| L_{n,\alpha}(|t - x|; x) \\ + \omega(f', \delta)(1 + \delta^{-1}\sqrt{L_{n,\alpha}((t - x)^2; x)})\sqrt{L_{n,\alpha}((t - x)^2; x)}. \end{aligned} \quad (5)$$

- (i) When $\alpha \geq 1$, taking $\delta = \frac{1}{\sqrt{n}}$ in (5), by Lemma 5 and inequality (5), we obtain the desired result.
- (ii) When $0 < \alpha < 1$, taking $\delta = n^{-\frac{\alpha}{2}}$ in (5), by Lemma 6 and inequality (5), we obtain the desired result. \square

Finally we study the direct theorem concerned with an approximation for the sequence $\{L_{n,\alpha}\}$ by means of the Ditzian-Totik modulus of smoothness. For the next theorem we shall use some notations.

For $f \in C[0, 1]$, $\varphi(x) = \sqrt{x(1-x)}$, $0 \leq \lambda \leq 1$, $x \in [0, 1]$, let

$$\omega_{\varphi^\lambda}(f, t) = \sup_{0 < h \leq t} \sup_{x \pm \frac{h\varphi^\lambda(x)}{2} \in [0, 1]} \left| f\left(x + \frac{h\varphi^\lambda(x)}{2}\right) - f\left(x - \frac{h\varphi^\lambda(x)}{2}\right) \right|$$

be the Ditzian-Totik modulus of first order, and let

$$K_{\varphi^\lambda}(f, t) = \inf_{g \in W_\lambda} \{ \|f - g\| + t \|\varphi^\lambda g'\| \} \quad (6)$$

be the corresponding K -functional, where $W_\lambda = \{f | f \in AC_{\text{loc}}[0, 1], \|\varphi^\lambda f'\| < \infty, \|f'\| < \infty\}$.

It is well known that (see [14])

$$K_{\varphi^\lambda}(f, t) \leq C\omega_{\varphi^\lambda}(f, t), \quad (7)$$

for some absolute constant $C > 0$.

Now we state our next main result.

Theorem 4 Let $f \in C[0, 1]$, $\alpha \geq 1$, $\varphi(x) = \sqrt{x(1-x)}$, $x \in [0, 1]$, $0 \leq \lambda \leq 1$. Then there exists an absolute constant $C > 0$ such that

$$|L_{n,\alpha}(f; x) - f(x)| \leq C\omega_{\varphi^\lambda}\left(f, \frac{\varphi^{1-\lambda}(x)}{\sqrt{n}}\right).$$

Proof Let $g \in W_\lambda$, by Lemma 2(i) and Lemma 7, we have

$$\begin{aligned} |L_{n,\alpha}(f; x) - f(x)| &\leq |L_{n,\alpha}(f - g; x)| + |f(x) - g(x)| + |L_{n,\alpha}(g; x) - g(x)| \\ &\leq 2\|f - g\| + |L_{n,\alpha}(g; x) - g(x)|. \end{aligned} \quad (8)$$

Since $g(t) = \int_x^t g'(u) du + g(x)$, $L_{n,\alpha}(1; x) = 1$, so, we have

$$\begin{aligned} |L_{n,\alpha}(g; x) - g(x)| &\leq \left| L_{n,\alpha}\left(\int_x^t |g'(u)| du; x\right) \right| \\ &\leq \|\varphi^\lambda g'\| L_{n,\alpha}\left(\left|\int_x^t \varphi^{-\lambda}(u) du\right|; x\right). \end{aligned} \quad (9)$$

By the Hölder inequality, we get

$$\left| \int_x^t \varphi^{-\lambda}(u) du \right| \leq \left| \int_x^t \frac{1}{\sqrt{u(1-u)}} du \right|^\lambda |t - x|^{1-\lambda}, \quad (10)$$

also, in view of $1 \leq \sqrt{u} + \sqrt{1-u} < 2$, $0 \leq u \leq 1$, we have

$$\begin{aligned} \left| \int_x^t \frac{1}{\sqrt{u(1-u)}} du \right| &\leq \left| \int_x^t \left(\frac{1}{\sqrt{u}} + \frac{1}{\sqrt{1-u}} \right) du \right| \\ &\leq 2(|\sqrt{t} - \sqrt{x}| + |\sqrt{1-x} - \sqrt{1-t}|) \\ &\leq 2\left(\frac{|t-x|}{\sqrt{t} + \sqrt{x}} + \frac{|t-x|}{\sqrt{1-t} + \sqrt{1-x}} \right) \\ &\leq 2|t-x| \left(\frac{1}{\sqrt{x}} + \frac{1}{\sqrt{1-x}} \right) \\ &\leq 4|t-x|\varphi^{-1}(x), \end{aligned} \quad (11)$$

thus, by (10) and (11), we obtain

$$\left| \int_x^t \varphi^{-\lambda}(u) du \right| \leq C\varphi^{-\lambda}(x)|t-x|, \quad (12)$$

also, by (9) and (12), we have

$$\begin{aligned} |L_{n,\alpha}(g; x) - g(x)| &\leq C\|\varphi^\lambda g'\| L_{n,\alpha}(\varphi^{-\lambda}(x)|t-x|; x) \\ &= C\|\varphi^\lambda g'\| \varphi^{-\lambda}(x) L_{n,\alpha}(|t-x|; x). \end{aligned} \quad (13)$$

In view of (4) and Lemma 2(i), by the Cauchy-Schwarz inequality, we have

$$\begin{aligned} L_{n,\alpha}(|t-x|;x) &\leq \sqrt{L_{n,\alpha}(1;x)}\sqrt{L_{n,\alpha}((t-x)^2;x)} \\ &\leq \sqrt{\frac{(n+1)\alpha}{n^2+1}}x(1-x) \\ &\leq C\frac{\varphi(x)}{\sqrt{n}}, \end{aligned} \quad (14)$$

so, by (13) and (14), we obtain

$$|L_{n,\alpha}(g;x) - g(x)| \leq C\|\varphi^\lambda g'\| \frac{\varphi^{1-\lambda}(x)}{\sqrt{n}}, \quad (15)$$

thus, by (8) and (15), we have

$$|L_{n,\alpha}(f;x) - f(x)| \leq 2\|f - g\| + C\|\varphi^\lambda g'\| \frac{\varphi^{1-\lambda}(x)}{\sqrt{n}}. \quad (16)$$

Then, in view of (16), (6), and (7), we obtain

$$|L_{n,\alpha}(f;x) - f(x)| \leq CK_{\varphi^\lambda}\left(f, \frac{\varphi^{1-\lambda}(x)}{\sqrt{n}}\right) \leq C\omega_{\varphi^\lambda}\left(f, \frac{\varphi^{1-\lambda}(x)}{\sqrt{n}}\right),$$

where C is a positive constant, in different places the value of C may be different. \square

4 Conclusions

In this paper, a new kind of type Bézier operators is introduced. The Korovkin type approximation theorem of these operators is investigated. The rates of convergence of these operators are studied by means of the modulus of continuity. Then, by using the Ditzian-Totik modulus of smoothness, a direct theorem concerned with an approximation for these operators is obtained. Further, we can also study the inverse theorem and an equivalent theorem concerned with an approximation for these operators.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Author details

¹School of Mathematics and Computer Science, Wuyi University, Wuyishan, 354300, China. ²School of Mathematical Sciences, Xiamen University, Xiamen, 361005, China.

Acknowledgements

This work is supported by the National Natural Science Foundation of China (Grant No. 61572020), the Class A Science and Technology Project of Education Department of Fujian Province of China (Grant No. JA12324), and the Natural Science Foundation of Fujian Province of China (Grant Nos. 2014J01021 and 2013J01017).

Received: 13 September 2015 Accepted: 9 December 2015 Published online: 23 December 2015

References

1. Bézier, P: Numerical Control: Mathematics and Applications. Wiley, London (1972)
2. Chang, GZ: Generalized Bernstein-Bézier polynomial. *J. Comput. Math.* **1**(4), 322-327 (1983)
3. Zeng, XM, Chen, W: On the rate of convergence of the generalized Durrmeyer type operators for functions of bounded variation. *J. Approx. Theory* **102**, 1-12 (2000)

4. Zeng, XM, Gupta, V: Rate of convergence of Baskakov-Bézier type operators for locally bounded functions. *Comput. Math. Appl.* **44**, 1445-1453 (2002)
5. Deo, N: Direct and inverse theorems for Szász-Lupas type operators in simultaneous approximation. *Mat. Vesn.* **58**, 19-29 (2006)
6. Guo, S, Qi, Q, Liu, G: The central approximation theorem for Baskakov-Bézier operators. *J. Approx. Theory* **147**, 112-124 (2007)
7. Liang, L, Sun, W: On pointwise approximation for a type of generalized Kantorovich-Bézier operators. *J. Southwest Univ.* **33**(10), 103-106 (2011)
8. Liu, G, Yang, X: On the approximation for generalized Szász-Durrmeyer type operators in the space $L_p[0, \infty)$. *J. Inequal. Appl.* **2014**, 447 (2014). doi:10.1186/1029-242X-2014-447
9. Deng, X, Wu, G: On approximation of Bernstein-Durrmeyer-Bézier operators in Orlicz spaces. *Pure Appl. Math.* **31**(3), 307-317 (2015)
10. Ren, MY: Approximation properties of a kind of Bernstein type operators. *J. Wuyi Univ.* **31**(2), 1-4 (2012)
11. Li, P, Huang, Y: Approximation order generalized Bernstein-Bézier polynomials. *J. Univ. Sci. Technol. China* **15**(1), 15-18 (1985)
12. Li, Z: Approximation properties of the Bernstein-Kantorovich-Bézier polynomials. *J. Nat. Sci. Hunan Norm. Univ.* **9**(1), 14-19 (1986)
13. Chen, WZ: *Operators Approximation Theory*. Xiamen University Press, Xiamen (1989) (in Chinese)
14. Ditzian, Z, Totik, V: *Moduli of Smoothness*. Springer, New York (1987)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com